Importance of post-donation information

Micheline Lambermont Service du Sang Belgian Red Cross

IHS 2013



Definition:

All information concerning the donor which

> becomes known only after the donation

> could have an implication on the quality and/or safety of one or more previous donations



Information sources :

- > Donor PDI : information transmitted directly by the donor himself
- > Testing PDI : information coming from the testing laboratory (sero-conversions...)
- Medical selection PDI : information detected at a subsequent donation



Donor PDI

- > 1996 : legally introduced in Belgium for autoexclusion (AIDS)
- > 2007 : enlarged to other donor information including delayed blood collection reaction
- > SFS : centralisation, unique telephone number, anonymous contact (donation number). Physician available 24h/24.



Donor PDI

Dans certaines situations, il existe un risque qu'une affection dont vous seriez atteint sans le savoir soit transmise à un malade.

- Si vous pensez qu'il y a un risque à transfuser votre sang suite aux informations qui vous ont été données...,
- Si vous avez oublié de signaler un élément important au médecin.
- Si, après votre don, vous développez une maladie infectieuse se traduisant par de la fièvre, une éruption cutanée, des ganglions ou des troubles digestifs. Vous vous trouviez peut-être, au moment de votre don, dans la période d'incubation de cette affection qui pourrait dès lors être transmise au receveur.

Dans l'un de ces cas, veuillez contacter IMMEDIATEMENT un médecin du Service du Sang au 078 051 053 (appel normal) en précisant le numéro du don repris sur cette carte... Pour toute autre info, appelez le 0800 92 245

SH6_POSDON_F_03

Service du Sang


Post-Don Nous vous remercions pour votre don de sang, Date de Naissance: Numéro Donneur:


Pour plus d'informations: www.transfusion.be 0800 92 245 - info@croix-rouge.be



Dete :	6 Date :
2	7
Date :	Date :
3 Date :	s Date :
4	9
Date :	Date :
5	10
Date :	Date :








Donor PDI Data 01/2007-12/2011 5 years

	Issued BC	Recalled BC	Transfused BC	Rate TBC
Total	245	158	87 (35%)	1/12.000
RBC	166	128	38 (23%)	1/20.000
PC	78	30	48 (61%)	1/2.000
VI FFP	2	1	1	<1/100.000


BC Distribution		BC Donor PDI rejection
CED :	803.651	1231
PC :	105.513	
VIFFP :	137.770	








Donor PDI delays







Issuing a blood component whether transfused (*incident*) or not (*near miss*) that does not fit the required quality and/or safety criteria

- > are the most numerous (48% of the total PDI)
- > are in majority (81%) due to infections
- > are reported with a mean delay of 4 days 42 % within 3 days, 75 % within a week
- > 1/3 leads to blood component rejection
- > 35% of BC already issued are transfused
- > 55 % (rate 1/2.000 issued plts) are platelet concentrates, pathogen reduced (Intercept Cerus) since may 2009, which is a main advantage for blood safety




Conclusion (2)

PDI : fundamental part of the hemovigilance system

PDI management implies :

- A very reactive organization
- an implication and sensitization of the BTE-SFS staff and especially the medical team
- A good awareness of the blood donors
- A close collaboration with the hospital blood bank responsible and clinicians







Thank you very much for your attention

