

POST-DONATION INFORMATION: AN IMPORTANT ELEMENT OF DONOR HAEMOVIGILANCE

Dra. T. Jimenez-Marco

Fundació Banc de Sang i Teixits de les Illes Balears, Majorca, Spain

POST-DONATION INFORMATION: AN IMPORTANT ELEMENT OF DONOR HAEMOVIGILANCE

Dr. T. Jimenez-Marco

Fundació Banc de Sang i Teixits de les Illes Balears, Majorca, Spain March 6, 2014

- The Balearic Island Hemovigilance Division (BIHVD) located in the Balearic Island Blood Bank (BIBB) facility, one of the 17 regional Hemovigilance divisions in the Spanish Hemovigilance network, collects and analyses all serious transfusion events reported by the Balearic Island Medical Centers..
- The BIHVD system, as well as the Spanish Hemovigilance Network, meet the requirement of the European Directive 2005/61/EC on Hemovigilance, and are based on an anonymous, voluntary, non-punitive serious adverse events reporting program.




Balearic Island Hemovigilance Division



- Early post-donation information (ePDI) is an essential element of donor haemovigilance since it helps to monitor the safety in donor screening process through medical interview.
- ePDI notification can be defined as deferrable medical history or donor behaviour not disclosed at the time of donation, but it is subsequently notified by the blood donors or by other reliable source before the following donation.

During the years 2005 and 2013, we analyzed the number and the categories of ePDI that were voluntary reported by the blood donors or by other reliable source early after the donation.






Early notification



Voluntary Blood Donor



BIHVD registration



- The ePDI were classified in the following categories:
 - Mild infections as flu, cold and gastrointestinal infections.
 - High-risk—sexual behaviour
 - High-risk—nonsexual behaviour as having close contact with a person viral hepatitis, having a tattoo, ear or skin piercing, acupuncture or coming into contact with someone else's blood
 - Travelling history to or being born in an endemic area either for parasite diseases as Malaria and Chagas disease or for other infectious disease such as HTLV I/II infection and Creutzfeldt-Jacob disease
 - A deferrable medical history as cancer or epilepsy, etc.
 - A deferrable medical treatment as finasteride, isotretinoin, etc. and
 - Blood diseases as bleeding disorders.

| Vacus | 2005 | 2007 | 2007 | 2000 | 2000 | 2010 | 2011 | 2012 | 2012 | Т.4.1 | D ((0() |
|--|--------|--------|--------|--------|--------|--------|--------|--------|--------|---------|----------------|
| Years | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | Total | Percentage (%) |
| Donations | 40,835 | 39,475 | 38,602 | 41,185 | 43,869 | 43,414 | 42,181 | 40,054 | 38,168 | 367,783 | |
| Donors | 21,563 | 22,54 | 22,486 | 25,006 | 27,722 | 26,121 | 25,294 | 24,491 | 22,839 | 218,062 | |
| ePDI | 23 | 29 | 20 | 17 | 27 | 20 | 29 | 23 | 26 | 214 | 100 |
| Mild Infections | 14 | 13 | 6 | 8 | 16 | 14 | 13 | 14 | 11 | 109 | 50.9 |
| High-risk- sexual | 2 | 5 | 3 | | 3 | 1 | 7 | 3 | 4 | 28 | 13.1 |
| High-risk- nonsexual | 5 | 5 | 7 | 4 | 5 | | 3 | 1 | 6 | 36 | 16.8 |
| Travel to or born in an endemic country | | 5 | 1 | 2 | 1 | 1 | 3 | 1 | 1 | 15 | 7.0 |
| Medical history | | | | 2 | | 3 | 2 | 1 | 2 | 10 | 4.7 |
| Medical treatment | 2 | 1 | 2 | 1 | | 1 | | | 0 | 7 | 3.3 |
| Blood disease | | | 1 | | 2 | | 1 | 3 | 2 | 9 | 4.2 |

- The most common ePDI notification is the mild infections symptoms reported by donors few days after giving blood.
- It is likely that donors donate blood during the asymptomatic incubation period of the infection.
- In order to prevent from using their donation for transfusion it is important to encourage donors to contact with the Blood Bank staff early after donation as soon as infection's symptoms appear.

- Other interesting aspect that needs to be investigated is the reason why some donors either forgot or intentionally fail to give information about their high-risk—nonsexual, high-risk—sexual behaviours and other conditions during the medical interview at the time of the donation.
- In our experience, the ePDI notification can be considered a very useful tool for monitoring medical interview accuracy.

Thank you very much



Questions?